

Supplementray file

Table S1. A list of primers used in this study

Note	Name	Primer sequences (5' to 3')
Real-time PCR	<i>EO</i> - qRT-F	CGTTTCGCAGCGGGTGTC
	<i>EO</i> - qRT-R	CCGATTTGGGTTTCAGCAGG
	<i>3DE-3β-reductase</i> - qRT-F	ATTCAAATAGAGGTTACCTC
	<i>3DE-3β-reductase</i> - qRT-R	GCGCTCAATCTGCCATCTCAAG
	<i>BmEck</i> -qRT-F	CGACCCGCTCACCGTCTACTC
	<i>BmEck</i> -qRT-R	GCTGCTCGTCTTTCACCTTG
	<i>RPL3</i> -qRT-F	CGGTGTTGTTGGATAATTGAG
	<i>RPL3</i> -qRT-R	GTCATCCTGCCATTTCTTACT
RNAi	ds-EGFP -F	^a <u>GTAATACGACTCACTATAGGGAGAGCTGTTACCGGGGTGGTGC</u>
	ds-EGFP -R	^a <u>GTAATACGACTCACTATAGGGAGATTCTCGTTGGGGTCTTTGCT</u>
	ds-EO-F	^a <u>TAATACGACTCACTATAGGCAGGATGTTTTTCGCACA</u>
	ds-EO-R	^a <u>TAATACGACTCACTATAGGAATTCGTATCAGCAATAG</u>
	ds-3DE-3β-reductase-F	^a <u>GTAATACGACTCACTATAGGGAGAAGAGGGTGCTGAAGGAAG</u>
	ds-3DE-3β-reductase-R	^a <u>GTAATACGACTCACTATAGGGAGAGGATTATCAACCTTGTCG</u>

^a The sites with wave line represent T7 promoter region.